




U.S. Environmental Protection Agency Integrated Risk Information System

[Recent Additions](#) | [Contact Us](#) | [Print Version](#)Search: [EPA Home](#) > [Browse EPA Topics](#) > [Human Health](#) > [Health Effects](#) > [IRIS Home](#) > [IRIS Summaries](#)

Benzo[g,h,i]perylene (CASRN 191-24-2)

[view QuickView](#)[List of IRIS Substances](#)Select a Substance 

go

☒ Full IRIS Summary ☐ QuickView

MAIN CONTENTS

[Reference Dose for Chronic Oral Exposure \(RfD\)](#)

0461

Benzo[g,h,i]perylene; CASRN 191-24-2

Health assessment information on a chemical substance is included in IRIS only after a comprehensive review of chronic toxicity data by U.S. EPA health scientists from several Program Offices and the Office of Research and Development. The summaries presented in Sections I and II represent a consensus reached in the review process. Background information and explanations of the methods used to derive the values given in IRIS are provided in the Background Documents.

STATUS OF DATA FOR Benzo[g,h,i]perylene

File First On-Line 12/01/1990

Category (section)	Status	Last Revised
Oral RfD Assessment (I.A.)	no data	
Inhalation RfC Assessment (I.B.)	no data	
Carcinogenicity Assessment (II.)	on-line	12/01/1990

_I. Chronic Health Hazard Assessments for Noncarcinogenic Effects

_I.A. Reference Dose for Chronic Oral Exposure (RfD)

Substance Name -- Benzo[g,h,i]perylene
CASRN -- 191-24-2

Not available at this time.

[Back to top](#)

_I.B. Reference Concentration for Chronic Inhalation Exposure (RfC)

SUBSTANCE SUMMARY INDEX

[Chronic Health
Hazards for Non-
Carcinogenic Effects](#)
[Reference Dose for
Chronic Oral Exposure
\(RfD\)](#)

[- Oral RfD Summary](#)
[- Principal and
Supporting Studies](#)
[- Uncertainty and
Modifying Factors](#)
[- Additional Studies/
Comments](#)
[- Confidence in the
Oral RfD](#)
[- EPA Documentation
and Review](#)

[Reference
Concentration for
Chronic Inhalation
Exposure \(RfC\)](#)

[- Inhalation RfC
Summary](#)
[- Principal and
Supporting Studies](#)
[- Uncertainty and
Modifying Factors](#)
[- Additional Studies/
Comments](#)
[- Confidence in the
Inhalation RfC](#)
[- EPA Documentation
and Review](#)

[Carcinogenicity
Assessment for
Lifetime Exposure](#)

[Evidence for Human
Carcinogenicity](#)

[- Weight-of-Evidence
Characterization](#)
[- Human
Carcinogenicity Data](#)
[- Animal
Carcinogenicity Data](#)
[- Supporting Data for
Carcinogenicity](#)



183293

Substance Name -- Benzo[g,h,i]perylene
CASRN -- 191-24-2

Not available at this time.

[Back to top](#)

[Quantitative Estimate
of Carcinogenic Risk
from Oral Exposure](#)

- [Summary of Risk
Estimates](#)
- [Dose-Response Data](#)
- [Additional Comments](#)
- [Discussion of
Confidence](#)

II. Carcinogenicity Assessment for Lifetime Exposure

Substance Name -- Benzo[g,h,i]perylene
CASRN -- 191-24-2
Last Revised -- 12/01/1990

[Quantitative Estimate
of Carcinogenic Risk
from Inhalation
Exposure](#)

- [Summary of Risk
Estimates](#)
- [Dose-Response Data](#)
- [Additional Comments](#)
- [Discussion of
Confidence](#)

[EPA Documentation,
Review and, Contacts](#)

- [Bibliography](#)
- [Revision History](#)
- [Synonyms](#)

Section II provides information on three aspects of the carcinogenic assessment for the substance in question; the weight-of-evidence judgment of the likelihood that the substance is a human carcinogen, and quantitative estimates of risk from oral exposure and from inhalation exposure. The quantitative risk estimates are presented in three ways. The slope factor is the result of application of a low-dose extrapolation procedure and is presented as the risk per (mg/kg)/day. The unit risk is the quantitative estimate in terms of either risk per ug/L drinking water or risk per ug/cu.m air breathed. The third form in which risk is presented is a drinking water or air concentration providing cancer risks of 1 in 10,000, 1 in 100,000 or 1 in 1,000,000. The rationale and methods used to develop the carcinogenicity information in IRIS are described in The Risk Assessment Guidelines of 1986 (EPA/600/8-87/045) and in the IRIS Background Document. IRIS summaries developed since the publication of EPA's more recent Proposed Guidelines for Carcinogen Risk Assessment also utilize those Guidelines where indicated (Federal Register 61(79):17960-18011, April 23, 1996). Users are referred to Section I of this IRIS file for information on long-term toxic effects other than carcinogenicity.

II.A. Evidence for Human Carcinogenicity

II.A.1. Weight-of-Evidence Characterization

Classification -- D; not classifiable as to human carcinogenicity

Basis -- Based on no human data and inadequate animal data from lung implant, skin-painting and subcutaneous injection bioassays.

II.A.2. Human Carcinogenicity Data

None.

II.A.3. Animal Carcinogenicity Data

Inadequate. Benzo[g,h,i]perylene appeared to increase lung epidermoid tumors when administered with trioctanonin in a lung implant study (Deutsch- Wenzel et al., 1983). Benzo[g,h,i]perylene was tested for complete carcinogenic activity and initiating activity in mouse skin painting assays and did not produce positive results in either type of assay (Wynder and Hoffmann, 1959; Hoffmann and Wynder, 1966; Muller, 1968; Van Duuren et al., 1973). Benzo[g,h,i]perylene did not induce tumor formation when injected subcutaneously (Muller, 1968) and was tested as a cocarcinogen with benzo(a)pyrene (Van Duuren et al., 1973; Van Duuren and Goldschmidt, 1976).

In a lifetime implant study, 3-month-old female Osborne-Mendel rats (34 to 35/group)

received a lung implant of benzo[g,h,i]perylene in 0.05 mL of a 1:1 (v:v) mixture of beeswax and trioctanoin (Deutsch-Wenzel et al., 1983). Rats received either 0.16 mg (0.65 mg/kg), 0.83 mg (3.4 mg/kg) or 4.15 mg (17 mg/kg). Controls consisted of an untreated group and a group receiving an implant of the vehicle. Median survival times (weeks) were: untreated controls, 118; vehicle implant controls, 104; 0.16 mg dose, 109; 0.83 mg dose, 114; and 4.15 mg dose, 106. Epidermoid carcinomas in the lung and thorax were observed at the following incidences: 0/35, 0/35, 0/35, 1/35 (3%), and 4/34 (12%) for the untreated controls, vehicle controls, low-, mid-, and high-dose groups, respectively. The apparent increased incidence of tumors was not statistically significant and no distant tumors were seen.

Benzo[g,h,i]perylene was tested as both a complete carcinogen and as a tumor initiator in female Ha/ICR/mil Swiss albino mice (Hoffmann and Wynder, 1966; Wynder and Hoffman, 1959). Two groups of 20 mice received dermal applications of 0.1 or 0.05% benzo[g,h,i]perylene 3 times/week for 1 year. The mice were observed for 3 additional months and then sacrificed. No tumors were observed in the low-dose group and a papilloma was observed in the high-dose group in the tenth month. In a second part of the study, benzo[g,h,i]perylene was applied as an initiator to 30 mice. Ten separate applications of 0.1% each were given over a 2-day period; beginning 28 days later 2.5% croton oil was applied 3 times/week for the remainder of the year. These mice were observed for 3 additional months; 2/27 surviving mice had developed papillomas in this group. A control group of 30 mice received applications of 2.5% croton oil (3 times/week) without an initiator; no tumor or survival data were reported.

The ability of benzo[g,h,i]perylene to act as a cocarcinogen in female ICR/HA mice when combined with benzo[a]pyrene was examined in a series of experiments (Van Duuren et al., 1973; Van Duuren and Goldschmidt, 1976). The mice (50/group) were treated by dermal application with 21 ug benzo[g,h,i]perylene and 5 ug benzo[a]pyrene (in combination) 3 times/week for 1 year. At the end of the experiment, 20/37 mice had developed papillomas and 17/37 had developed squamous cell carcinomas. In the control group, which consisted of three benzo[a]pyrene treatments (5 ug/week), 13/42 mice developed papillomas and 10/42 developed carcinomas (Van Duuren et al., 1973). In the second experiment two doses of benzo[g,h,i]perylene (7 and 21 ug) were applied along with 5 ug benzo[a]pyrene to groups of 50 mice 3 times/week for 368 days. In the low-dose group 19 mice developed papillomas and 10 carcinomas; in the high-dose group, 20 mice developed papillomas and 18 carcinomas. No papillomas or carcinomas developed when 21 ug benzo[g,h,i]perylene was applied alone. The individual animal data were not given for this experiment.

In a series of experiments, Muller (1968) investigated the carcinogenicity of benzo[g,h,i]perylene. In the first experiment groups of 50 NMRI mice (sex unspecified) received dermal applications (2 or 3 times/week) of one of a variety of concentrations of benzo[g,h,i]perylene in dichloromethane. A control group receiving only 0.2 mL dichloromethane was also utilized. The study was terminated 675 days after the first application. Survival was approximately the same in all four groups (33%). No skin papillomas or carcinomas developed; however, both benign (0/18 low-, 2/14 mid- and 3/17 high-dose groups) and malignant (3/18 low-, 4/14 mid- and 1/17 high-dose groups) tumors in survivors did occur at other sites (types and sites not specified). In the control group, 3/17 mice developed benign tumors and 4/17 developed malignant tumors at other sites. Dichloromethane is classified B2, a probable human carcinogen.

In a second dermal application study, groups of 50 mice initially were untreated or treated with a single application of either 1 or 2 mg benzo[g,h,i]perylene. In each group repeated dermal applications of 0.2 mL of 0.5% croton oil (2 times/week) followed for 25 weeks. One mouse in the promoter control group and another in the high-dose group developed skin papillomas; 2/28 (0/28), 4/12 (1/12), and 2/21 (1/21) mice developed benign (malignant) tumors at other sites (unspecified) in the control,

low- and high- dose groups, respectively. In the third part of the experiment three groups of 50 female NMRI mice received subcutaneous injections of 0 (control), 0.83 or 16.7 mg benzo[g,h,i]perylene suspended in 0.15 mL 10% aqueous gelatin once every 2 weeks for 6 months [total doses, 0, 10 and 200 mg/animal] and observed to sacrifice on day 675 after the first injection. At that time the survival rate was 36% in each group. No tumor was found at the site of injection in any of the animals. For the control, low- and high-dose groups, respectively, 4/50, 5/50, and 4/50 mice had tumors at other sites. In the final part of the experiment four groups of 20 NMRI mice (sex unspecified) were given subcutaneous injections of 0.15 mL 10% aqueous gelatin containing 0 (control), 0.1, 1, or 10 mg suspended benzo[g,h,i]perylene (total doses, 0, 10, or 100 mg/animal) once every 2 weeks for 20 weeks. The animals were observed until spontaneous death. Survival was not adversely affected by treatment with benzo[g,h,i]perylene (the last animal died 22 months after the start of the study). There is no information to indicate if enough animals survived long enough for tumors to be seen. No skin or subcutaneous tumors were found in mice treated with benzo[g,h,i]perylene or gelatin. Few tumors were found in other organs and the incidences in the benzo[g,h,i]perylene-treated groups were not different from those in the gelatin controls. Earlier skin-painting studies are summarized in IARC (1983).

II.A.4. Supporting Data for Carcinogenicity

Benzo[g,h,i]perylene produced positive results in tests for reverse mutation in three strains of *Salmonella typhimurium* and for forward mutation in one strain (Andrews et al., 1978; Mossanda et al., 1979; Salamone et al., 1979; Sakai et al., 1985; Kaden et al., 1979). A test for DNA damage in Chinese hamster ovary cells also yielded positive results (Garrett and Lewtas, 1983).

[Back to top](#)

II.B. Quantitative Estimate of Carcinogenic Risk from Oral Exposure

None.

[Back to top](#)

II.C. Quantitative Estimate of Carcinogenic Risk from Inhalation Exposure

None.

[Back to top](#)

II.D. EPA Documentation, Review, and Contacts (Carcinogenicity Assessment)

II.D.1. EPA Documentation

Source Document -- U.S. EPA, 1990

The 1990 Drinking Water Criteria Document for Polycyclic Aromatic Hydrocarbons

has received Agency and external review.

__II.D.2. EPA Review (Carcinogenicity Assessment)

Agency Work Group Review -- 02/07/1990

Verification Date -- 02/07/1990

__II.D.3. EPA Contacts (Carcinogenicity Assessment)

Please contact the IRIS Hotline for all questions concerning this assessment or IRIS, in general, at (202)566-1676 (phone), (202)566-1749 (FAX) or hotline.iris@epa.gov (internet address).

[Back to top](#)

_III. [reserved]

_IV. [reserved]

_V. [reserved]

_VI. Bibliography

Substance Name -- Benzo[g,h,i]perylene

CASRN -- 191-24-2

Last Revised -- 12/01/1990

_VI.A. Oral RfD References

None

[Back to top](#)

_VI.B. Inhalation RfC References

None

[Back to top](#)

_VI.C. Carcinogenicity Assessment References

Andrews, A.W., L.H. Thibault and W. Lijinsky. 1978. The relationship between carcinogenicity and mutagenicity of some polynuclear hydrocarbons. *Mutat. Res.* 51: 311-318.

Deutsch-Wenzel, R., H. Brune, G. Grimmer, G. Dettbarn and J. Misfeld. 1983. Experimental studies in rat lungs on the carcinogenicity and dose-response

relationships of eight frequently occurring environmental polycyclic aromatic hydrocarbons. *J. Natl. Cancer Inst.* 71(3): 539-544.

Garrett, N.E. and J. Lewtas. 1983. Cellular toxicity in Chinese hamster ovary cell cultures. 1. Analysis of cytotoxicity endpoints for twenty-nine priority pollutants. *Environ. Res.* 32(2): 455-465.

Hoffmann, D. and E.L. Wynder, 1966. Beitrag zur carcinogen Wirkung von Dibenzopyrene. *Z. Krebsforsch.* 68(2): 137-149. (Ger.) Contribution on the carcinogenic effect of dibenzopyrenes.

IARC (International Agency for Research on Cancer). 1983. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals for Humans. Polynuclear Aromatic Compounds. Part 1. Chemical, Environmental and Experimental Data. Vol. 32. World Health Organization.

Kaden, D.A., R.A. Hites and W.G. Thilly. 1979. Mutagenicity of soot and associated polycyclic aromatic hydrocarbons to *Salmonella typhimurium*. *Cancer Res.* 39: 4152-4159.

Mossanda, K., F. Poncelet, A. Fouassin and M. Mercier. 1979. Detection of mutagenic polycyclic aromatic hydrocarbons in African smoked fish. *Food Cosmet. Toxicol.* 17: 141-143.

Muller, E. 1968. Carcinogenic substances in water and soil. XX. Studies on the carcinogenic properties of 1,12-benzoperylene. *Arch. Hyg. Bakteriolog.* 152: 23-36. (Ger.)

Sakai, M., D. Yoshida and S. Mizusaki. 1985. Mutagenicity of polycyclic aromatic hydrocarbons and quinones on *Salmonella typhimurium* TA97. *Mutat. Res.* 156: 61-67.

Salamone, M.F., J.A. Heddle and M. Katz. 1979. The mutagenic activity of thirty polycyclic aromatic hydrocarbons (PAH) and oxides on urban airborne particulates. *Environ. Int.* 2: 37-43.

U.S. EPA. 1990. Drinking Water Criteria Document for Polycyclic Aromatic Hydrocarbons (PAHs). Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC. Final Draft. ECAO-CIN-D010, September, 1990.

Van Duuren, B.L. and B.M. Goldschmidt. 1976. Cocarcinogenic and tumor-promoting agents in tobacco carcinogenesis. *J. Natl. Cancer Inst.* 56(6): 1237-1242.

Van Duuren, B.L., C. Katz and B.M. Goldschmidt. 1973. Brief communication: Cocarcinogenic agents in tobacco carcinogenesis. *J. Natl. Cancer Inst.* 51(2): 703-705.

Wynder, E.L. and D. Hoffmann. 1959. A study of tobacco carcinogenesis. VII. The role of higher polycyclic hydrocarbons. *Cancer.* 12: 1079-1086.

[Back to top](#)

_VII. Revision History

Substance Name -- Benzo[g,h,i]perylene
CASRN -- 191-24-2

Date	Section	Description
12/01/1990	II.	Carcinogen assessment on-line
12/01/1990	VI.	Bibliography on-line
01/01/1992	IV.	Regulatory Action section on-line
04/01/1997	III., IV., V.	Drinking Water Health Advisories, EPA Regulatory Actions, and Supplementary Data were removed from IRIS on or before April 1997. IRIS users were directed to the appropriate EPA Program Offices for this information.

[Back to top](#)

_VIII. Synonyms

Substance Name -- Benzo[g,h,i]perylene
CASRN -- 191-24-2
Last Revised -- 12/01/1990

191-24-2
Benzo(ghi)perylene
benzo(ghi)perylene
HSDB 6177
NSC 89275
1,12-Benzoperylene
1,12-benzperylene

[Back to top](#)

[Recent Additions](#) | [Search IRIS](#) | [IRIS Home](#) | [NCEA Home](#) | [ORD Home](#)

[EPA Home](#) | [Privacy and Security Notice](#) | [Contact Us](#)

Last updated on Thursday, November 18th, 2004
URL: <http://www.epa.gov/iris/subst/0461.htm>